Instructions for Use

25ga I.D.D. Internal Delivery Device (VS0225.25)

Indications for use - Vortex Surgical 25ga I.D.D. Internal Delivery Device is intended to inject, irrigate, and/or remove fluids during vitreoretinal surgery. Contraindications - None

Device Description - Vortex Surgical 25ga I.D.D. Internal Delivery Device has a curved nitinol needle that can be retracted within 25ga stainless steel needle. The I.D.D. is compatible with its corresponding gauge trocar cannula system.

Known complications - Retinal Detachment, tears, holes, contusions, bleeding, inflammation, infection, and partial loss of vision.

These complications are statistically rare and it is assumed that the user is adequately trained in the treatment of these known complications and methods of avoidance.

Caution - Federal (USA) law restricts this device sale by, or on the order of, a physician.

Instructions for use: (Electronic Copy of IFU can be found at https://www.vortexsurgical.com/instructionsforuse)

1. Determine package integrity. 2. Affect sterile transfer of the product to the sterile field. 3. Remove the instrument from the tray. 4. Visually inspect tip surface. Do not use if damaged. 5. The instrument is now prepared for use, 6. Follow established surgical procedures.

Warnings - The device is designed to be utilized with cannulas. Ensure the curved portion of the device is fully retracted prior to insertion through the valved or non-valved cannulas. Do not use if package integrity or product compromised. Do not use if product exposed to conditions outside of indicated range. Do not use if EtO indicator does not show exposure to sterilization gas.

<u>Device Storage</u> - Vortex Surgical, Inc. recommends that the product is stored in a clean, dry and well-ventilated area at room temperature 15-37 °C (59-98.6 °F) away from direct sunlight.

Sterilization - This product is sterilized by Ethylene Oxide Gas (EtO) and provided with an EtO indicator. Verify indicator status color prior to use. Re-sterilization - Do not re-sterilize instrument. Reuse - Do not reuse or reprocess instrument. Reuse could lead to cross contamination, infection, or device failure.

Device Disposal - This single use surgical instrument should be considered clinical waste and should be disposed in accordance with clinical waste laws applicable in your country.

Reporting - Report any serious incident that has occurred in relation to the device to Vortex Surgical and your regulatory body including the competent authority of the Member State in which the user and/or patient is established.

Vortex Surgical, Inc. excludes all warranties, whether expressed, implied, or otherwise, to matters beyond the direct control of Vortex Surgical, Inc. and the end result of this device's use. This would include, but not be limited to handling, shipment and storage of the device and patient diagnosis and treatment. The fitness and merchantability of this device are as specified. Implied factors are specifically excluded. Vortex Surgical, Inc. is not liable for loss, whether incidental or consequential, damage and/or expense, arising wither directly or indirectly from the use of this device. There is no additional liability or responsibility assumed, other than that specified. All additional liability or responsibility is specifically disclaimed.



Vortex Surgical, Inc., 4 Research Park Drive, Suite 124, St. Charles, MO 63304, USA info@vortexsurgical.com 636-778-4350 www.vortexsurgical.com





Authorized representative in the European Union

Do not







STERILE EO

Sterilized using

ethylene oxide



Temperature

limit

15°C

37° C



2200009 rev C

Instructions for Use

25ga I.D.D. Internal Delivery Device (VS0225.25)

Indications for use - Vortex Surgical 25ga I.D.D. Internal Delivery Device is intended to inject, irrigate, and/or remove fluids during vitreoretinal surgery. Contraindications - None

Device Description - Vortex Surgical 25ga I.D.D. Internal Delivery Device has a curved nitinol needle that can be retracted within 25ga stainless steel needle. The I.D.D. is compatible with its corresponding gauge trocar cannula system.

Known complications - Retinal Detachment, tears, holes, contusions, bleeding, inflammation, infection, and partial loss of vision.

These complications are statistically rare and it is assumed that the user is adequately trained in the treatment of these known complications and methods of avoidance.

Caution - Federal (USA) law restricts this device sale by, or on the order of, a physician.

Instructions for use: (Electronic Copy of IFU can be found at https://www.vortexsurgical.com/instructionsforuse)

1. Determine package integrity. 2. Affect sterile transfer of the product to the sterile field. 3. Remove the instrument from the tray. 4. Visually inspect tip surface. Do not use if damaged. 5. The instrument is now prepared for use. 6.Follow established surgical procedures.

Warnings - The device is designed to be utilized with cannulas. Ensure the curved portion of the device is fully retracted prior to insertion through the valved or non-valved cannulas. Do not use if package integrity or product compromised. Do not use if product exposed to conditions outside of indicated range. Do not use if EtO indicator does not show exposure to sterilization gas.

Device Storage - Vortex Surgical, Inc. recommends that the product is stored in a clean, dry and well-ventilated area at room temperature 15-37 °C (59-98.6 °F) away from direct sunlight.

Sterilization - This product is sterilized by Ethylene Oxide Gas (EtO) and provided with an EtO indicator. Verify indicator status color prior to use. Re-sterilization - Do not re-sterilize instrument. Reuse - Do not reuse or reprocess instrument. Reuse could lead to cross contamination, infection, or device failure.

Device Disposal - This single use surgical instrument should be considered clinical waste and should be disposed in accordance with clinical waste laws applicable in your country.

Reporting - Report any serious incident that has occurred in relation to the device to Vortex Surgical and your regulatory body including the competent authority of the Member State in which the user and/or patient is established.

Vortex Surgical, Inc. excludes all warranties, whether expressed, implied, or otherwise, to matters beyond the direct control of Vortex Surgical, Inc. and the end result of this device's use. This would include, but not be limited to handling, shipment and storage of the device and patient diagnosis and treatment. The fitness and merchantability of this device are as specified. Implied factors are specifically excluded. Vortex Surgical, Inc. is not liable for loss, whether incidental or consequential, damage and/or expense, arising wither directly or indirectly from the use of this device. There is no additional liability or responsibility assumed, other than that specified. All additional liability or responsibility is specifically disclaimed.



Vortex Surgical, Inc., 4 Research Park Drive, Suite 124, St. Charles, MO 63304, USA info@vortexsurgical.com 636-778-4350 www.vortexsurgical.com

Manufacturer

Keep

drv

Consult

Instructions

for Use



Authorized representative







if damaged





use only

STERILE



Sterilized using

ethylene oxide

EO







_37°C

C Vortex Surgical Inc., 2024

Issue Date: 2024-12-06

2200009 rev C



for Use

Keep

Instructions

Lot number

Keep away

from sunlight

LOT

REF

Do not

resterilize

number

Quantity Catalog

Use Βv

Do not used

if damaged

Manufactured in USA/Date of Manufacture

V√V USA

use only



Device

Not made with natural rubber

latex

LOT

number

Keep away

from sunlight

Lot

REF

Catalog

Do not

resterilize

number

QT' Quantity

Do not

reuse

Use Ву

Manufactured in USA/Date of Manufacture

/W USA

(L'ATEX Not made with natural rubber

Device

latex